



BLASTER* PRO



HERBICIDE

Product Registration Number: MAPP 15752

An emulsifiable concentrate containing
240 g/litre triclopyr (present as 334 g/litre of triclopyr butotyl) and 60 g/litre clopyralid.

A foliar acting herbicide for the control of NETTLES, DOCKS, THISTLES, BRAMBLES,
BROOM and GORSE in AMENITY GRASSLAND.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use
of this product at work.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET.

PROTECT FROM FROST.

0.25-5.0 Litre(s)

Dow AgroSciences Limited

Latchmore Court, Brand Street, Hitchin,
Hertfordshire. SG5 1NH.

Telephone: Hitchin (01462) 457272

Fax: (01462) 426605

24 Hour Emergency Telephone Number: +44 (0) 1553 761 251

Headland Amenity Ltd.

1010 Cambourne Business Park,
Cambourne,

Cambridgeshire CB3 6DP

Telephone: 01223 597834

*Trademark of Dow AgroSciences LLC.

This label is compliant with the CPA Voluntary Initiative Guidance.



SAFETY PRECAUTIONS

Operator protection:

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

- a) Operators must wear suitable protective clothing (coveralls), suitable protective gloves, rubber boots and face protection (faceshield) when handling the concentrate.
 - b) Operators must wear suitable protective clothing (coveralls), suitable protective gloves and rubber boots when applying by hand-held equipment.
 - c) Operators must wear suitable protective gloves when handling contaminated surfaces.
- However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

DO NOT APPLY by hand-held rotary atomiser equipment.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

WASH CONCENTRATE from skin or eyes immediately.

DO NOT BREATHE SPRAY.

WASH HANDS AND EXPOSED SKIN before meals and after work.

Consumer protection:

NOT TO BE USED ON FOOD CROPS.

Environmental protection:

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. **DO NOT ALLOW DIRECT SPRAY** from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

'Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area'.

DO NOT CONTAMINATE WATER with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

KEEP LIVESTOCK out of treated areas for at least 7 days and until foliage of any poisonous weeds such as ragwort has died and become unpalatable.

Storage and disposal:

KEEP OUT OF REACH OF CHILDREN.



KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.



HARMFUL



DANGEROUS FOR THE ENVIRONMENT

**HARMFUL: MAY CAUSE LUNG DAMAGE IF SWALLOWED.
IRRITATING TO EYES, RESPIRATORY SYSTEM AND SKIN.
MAY CAUSE SENSITISATION BY SKIN CONTACT.
VAPOURS MAY CAUSE DROWSINESS AND DIZZINESS.
TOXIC TO AQUATIC ORGANISMS, MAY CAUSE LONG-TERM ADVERSE EFFECTS IN THE AQUATIC ENVIRONMENT. WEAR SUITABLE GLOVES.
DO NOT BREATHE VAPOUR
WEAR SUITABLE GLOVES.
AVOID CONTACT WITH SKIN AND EYES
IN CASE OF CONTACT WITH EYES, RINSE IMMEDIATELY WITH PLENTY OF WATER AND SEEK MEDICAL ADVICE.
IF SWALLOWED, DO NOT INDUCE VOMITING: SEEK MEDICAL ADVICE IMMEDIATELY AND SHOW THIS CONTAINER OR LABEL.
THIS MATERIAL AND ITS CONTAINER MUST BE DISPOSED OF IN A SAFE WAY. USE APPROPRIATE CONTAINMENT TO AVOID ENVIRONMENTAL CONTAMINATION.
To avoid risks to man and the environment, comply with the instructions for use.**

IMPORTANT INFORMATION.

FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops/Situations: Amenity grassland
Maximum Individual Dose: 1.2 litres product per hectare (see Other Specific Restrictions)
Maximum Number of Treatments: One per year

Other Specific Restrictions:

DO NOT APPLY through hand-held rotary atomiser equipment.
The maximum concentration must not exceed 60 mL product per 10 litres of water (6 mL product per litre of water)
KEEP LIVESTOCK out of treated areas for at least 7 days and until foliage of any poisonous weeds such as ragwort has died and become unpalatable.
Applications must not be made outside the period of 1st March to 31st October

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

WARNINGS

Read all warnings before applying BLASTER PRO.

Do not roll or harrow grass for 10 days before or 7 days after application.

Do not spray in drought, very hot or very cold weather.

BLASTER PRO is safe to grass. Very occasionally some yellowing of the sward may occur; this is transient and quickly outgrown.

Clover will be killed or severely checked by application of BLASTER PRO.

DO NOT sow kale, swedes, turnips or grass mixtures containing clover by direct drilling or minimum cultivation techniques within 6 weeks of applying BLASTER PRO.

BLASTER PRO residues in plant tissues (including manure and digestate) which have not completely decayed may affect succeeding susceptible crops e.g. peas, beans and other legumes, carrots and Umbelliferae, potatoes and tomatoes, lettuce and other Compositae. Do not plant susceptible autumn-sown crops (e.g. winter beans) in the same year as treatment with BLASTER PRO. Where susceptible crops are to be planted in the spring do not apply BLASTER PRO later than the end of July of the previous year. Following good agricultural practice ensure that plant remains have completely decayed before planting susceptible crops.

Do not use any plant material treated with BLASTER PRO for composting or mulching.

Do not use manure from animals fed on crops treated with BLASTER PRO for composting.

In hot conditions, vapour drift may occur, making it particularly important to assess the risk to neighbouring vegetation.

Take extreme care to avoid drift onto crops and non-target plants outside the target area.

Do not apply directly to, or allow spray drift to come into contact with agricultural or horticultural crops, amenity plantings or gardens, ponds, lakes or water courses.

All conifers, especially pine and larch, are very sensitive to spray drift.

Grazing Interval

Exclude livestock during treatment and do not allow livestock to graze treated grassland for at least 7 days following treatment, and until foliage of any poisonous weeds which may have been affected by application has died and become unpalatable.

WEEDS CONTROLLED, RATE OF USE AND APPLICATION TIMING

The following weeds will be controlled by an application of BLASTER PRO at a rate of 60 mL of product in 10 litres of water (6 mL product in 1 litre of water).

The weeds should be thoroughly wetted with the spray solution but spraying until "run-off" will decrease activity. The use of flood jets is recommended to prevent drift. Care should be taken to avoid local overdosing.

If the grass has been cut for hay or silage or grazed leave for 2-3 weeks to allow sufficient re-growth to occur before spraying. Grass and weeds must be actively growing to ensure good weed control and minimal check to the grass.

To allow maximum translocation of BLASTER PRO to the roots do not cut grass for 28 days after application.

Weeds	Optimum timing of application
Common nettle	Spray when actively growing but preferably before flowering (normally up to mid-June).
Curled dock Broad-leaved dock	Treat in the spring when the docks are in the rosette stage up to 25 cm high. On large well-established docks, or where there is a high reservoir of seed in the soil, a second application the following year may be required.
Creeping thistle	Spray when actively growing but before flowering spikes are 15 cm high. Application of BLASTER PRO at flowering or during seeding is likely to produce reduced levels of control.
Bramble, broom and gorse	Spray in June-August when actively growing but before plants begin to senesce in the autumn. It is essential that, particularly with large bushes, all the foliage is thoroughly wetted or incomplete kill may result.

APPLICATION EQUIPMENT

Apply BLASTER PRO using a handheld sprayer. Ensure the equipment is in good working order and has been calibrated according to the manufacturers' recommendations.

Wash out spray equipment thoroughly with water and detergent immediately after use. Traces of BLASTER PRO could cause harm to susceptible crops sprayed later.

DO NOT APPLY through hand-held rotary atomiser equipment.

Dow AgroSciences Conditions of Supply

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use, or the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded. No responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

[INSERT SDS HERE]